



NGOC Clinical Trials List - Printable Version

Head and Neck (C00-C14)

1. Locally Advanced
2. 1st Line Recurrent / Metastatic
3. Previously Treated

Esophagus/Gastric (C15-C16)

1. Stage II- IV, Neo-adjuvant
2. First Line, Recurrent or Metastatic
 - a. **AZ D702AC00001 (Artemide-Gastric01) A Phase 3 Study of Rilvegostomig in Combination with Fluoropyrimidine**
 - i. Arm A: Rilvegostomig + T-Dxd + Investigator's Choice: 5-FU or Capecitabine
 - ii. Arm B: Pembro + trastuzumab + Investigator's choice: Cisplatin+ 5FU or CAPOX+capecitabine
 - iii. Arm C: Rilvegostomic (750mg IV Q3w) + Trastuzumab + Investigator's Choice: Cisplatin+5FU OR CAPOX+Capecitabine
3. Previously Treated

Open 8/12/25

Biliary (K83.9)

1. First Line Therapy
 - a. **AZ D702NC00001 A Global Phase III Study of Rilvegostomig Plus Chemotherapy for First-**
 - i. Arm A: Rilvegostomig 750mg on D1 of each Q3W+ Gemcitabine 1000 mg/m2 IV Q3W plus cisplatin 25 mg/m2 IV Q3W will be administered on D1 and D8 of each cycle starting on C1D1 for up to 8 cycles
 - ii. Arm B:Durvalumab 1500mg on D1 of each Q4W cycle +Gemcitabine 1000 mg/m2 IV Q3W plus cisplatin 25 mg/m2 IV Q3W will be administered on D1 and D8 of each cycle starting on C1D1 for up to 8 cycles

Open 3/25/26

Pancreas (C25)

1. Adjuvant / NeoAdjuvant
2. 1st Line, Metastatic
 - a. **BMS CA240-0030 (MountainTAP) Study of BMS-986504 in Combination with Nab-p/Gem versus Placebo in Co**
 - i. CLOSED: Phase 2 Dose Selection Arm A: BMS-986504 400mg + Nab-Pac + Gem
 - ii. CLOSED: Phase 2 Dose Selection Arm B: BMS-986504 600mg + Nab-Pac + Gem
 - iii. CLOSED: Phase 2 Dose Selection Arm C: BMS-986504 400mg Placebo + Nab-Pac + Gem
 - iv. CLOSED: Phase 2 Dose Selection Arm D: BMS-986504 600mg Placebo + Nab-Pac + Gem
 - v. NOT YET OPEN Phase 3 Arm E: BMS-986504 Optimal Dose + Nab-Pac + Gem
 - vi. NOT YET OPEN Phase 3 Arm F: BMS-986504 Optimal Dose Placebo + Nab-Pac + Gem
3. Previously Treated
 - a. **RevMed RMC-6236-401 EXPANDED ACCESS PROGRAM TO TREAT PATIENTS WITH PREVIOUSLY TREATED MET**
 - i. Single Arm: Daraxonrasib 300mg PO QD for 28d Cycle

On Hold 5/5/26

Open 7/14/26

Colon & Rectum (C18-C20)

1. Stage III, Resected Colon, Adjuvant
2. Stage IV, Metastatic, 1st Line, Colon or Rectum
3. Metastatic, 1st Line, Colorectal
 - a. **BMS CA266-0003 (ROSETTA CRC-203) Punitamig in Combination with Chemotherapy Versus Bevacizumab in Combi**
 - i. Closed-Phase 2 Arm A1: punitamig (BMS-986545) Dose Level 1, Day 1 + FOLFOX or FOLFIRI,Q2W
 - ii. Closed-Phase 2 Arm A2: punitamig (BMS-986545) Dose Level 2, Day 1 + FOLFOX or FOLFIRI,Q2W
 - iii. Closed-Phase 2 Arm B:bevacizumab, Day 1 + FOLFOX or FOLFIRI, Q2W
 - iv. NOT OPEN Phase 3 Arm C: punitamig RP3D, Day 1 + FOLFOX or FOLFIRI, Q2W; or punitamig RP3D + CAPOX, Q3W
 - v. NOT OPEN Phase 3 Arm D:bevacizumab, Day 1 + FOLFOX or FOLFIRI, Q2W or bevacizumab + CAPOX, Q3W
 - b. **AbbV M24-533 (AndroMETA-CRC-533) A Phase 2, Open-Label, Randomized, Master Protocol Study to Evaluate S**
 - i. PENDING Stage 2 - SubStudy 1 Arm A: Folfox+Bevacizumab+Telisotuzumab Adizutecan High dose (2.4mg/kg)
 - ii. PENDING Stage 2 - SubStudy 1 Arm B: Folfox+Bevacizumab+Telisotuzumab Adizutecan Low dose (2.0mg/kg)
 - iii. PENDING Stage 2 - SubStudy 1 Arm C: SOC: Folfox+Bevacizumab
 - iv. Stage 2 - SubStudy 2 Arm A: 5FU/leucovorin+Panitumumab+Telisotuzumab Adizutecan High dose (2.4mg/kg)
 - v. Stage 2 - SubStudy 2 Arm B: 5FU/leucovorin+Panitumumab+Telisotuzumab Adizutecan Low dose (2.0mg/kg)
 - vi. Stage 2 - SubStudy 2 Arm C: FOLFOX+Panitumumab
4. Metastatic, 3rd Line, Colon
5. Metastatic 2nd or 3rd Line, KRAS Mutated
6. Metastatic 2nd or 3rd Line, KRAS Wild Type
7. Refractory

Pending 4/28/26

Pending 6/12/26

Lung, Squamous Cell (C34)

1. Stage IV, Metastatic
2. Second Line

Lung, Non-Small Cell (C34)

1. Stage IB, II, IIIA; Adjuvant
2. Stage 1B, II, Resectable IIIA
3. Stage II-III Resectable or Resected

a. **BI 1479-0032 (Beamion Lung-3) A Phase 3, Beamion LUNG-3: A study to test whether zongertinib helps**

- i. Arm A: Zongertinib (BI 1810631) 120mg po qd
- ii. Arm B: Physicians Choice(Nivolumab 480mg, pembrolizumab 200mg, atezolizumab 1200mg, or durvalumab 1500mg)

Open 3/19/26

4. Stage III, Locally Advanced, Unresectable

a. **BMS CA266-0001 (ROSETTA Lung-201) Study of Punitamig Vs Durvalumab Following Concurrent Chemoradiation T**

- i. Arm A: Punitamig 1200 mg (if body weight < 50 kg) or 1500 mg (if body weight ≥ 50 kg) 60-minute IV Q3W
- ii. Arm B: Durvalumab 1500 mg 60-minute IV Q4W

Open 7/22/26

5. Stage IIB-IIIIB Resected and Chemotherapy-Naive Stage IV

6. Stage IV, Metastatic

7. First Line- Metastatic

a. **AZ D702GC00001 Artemide Lung-04 A Global Phase III Study of Rilvegostomig or Pembrolizumab Monotherapy**

- i. Arm A: Rilvegostomig 750 mg IV Q3W
- ii. ArmB: Pembrolizumab 200 mg IV Q3W

Open 1/7/26

b. **BMS CA266-0002(ROSETTA Lung-202) A Study of Punitamig Versus Pembrolizumab in Participants With Previou**

- i. Arm A: Punitamig 1200 mg (if body weight < 50 kg) or 1500 mg (if body weight ≥ 50 kg) as a 60-minute IV infusion every 3 weeks (Q3W)
- ii. Arm B: Pembrolizumab 200 mg as a 60-minute IV infusion every 3 weeks (Q3W)

Open 4/15/26

c. **BMS CA239-0004 (KRYSTAL-4) Adagrasib plus Pembrolizumab plus Chemotherapy vs. Placebo plus Pembro**

- i. Arm A: Adagrasib 400mg BID + pemetrexed 500mg/m2 Q3W (until PD or unacceptable toxicity) + [cisplatin 75mg/m2 OR carboplatin AUC 5mg/mL/min Q3W (4 cycles)] + pembrolizumab 200mg Q3W (up to 24 months)
- ii. Arm B: Placebo + pemetrexed 500 mg/m2 Q3W (until PD or unacceptable toxicity) + [cisplatin 75 mg/m2 OR carboplatin AUC 5 mg/mL/min Q3W (4 cycles)] + pembrolizumab 200 mg Q3W (up to 24 months)

Open 5/19/26

8. 2nd or 3rd Line

9. Metastatic, Previously Treated

a. **AbbV M25-274 (TEImet NSCLC-04) A Phase 2, Open-Label, Randomized, Global Study of Three Telisotuzumab**

- i. Arm A: Telisotuzumab vedotin 1.9 mg/kg Q2W
- ii. Arm B: Telisotuzumab vedotin 1.6 mg/kg Q2W
- iii. Arm C: Telisotuzumab vedotin 1.6 mg/kg Q2W x 4 cycles then 1.9 mg/kg Q2W

Open 10/20/25

b. **AZ D763QC00001 (TROPION-Lung17) Datopotamab Deruxtecan or Docetaxel in Previously Treated TROP2-posit**

- i. Arm A: Dato-DXd 6mg/kg IV q3w
- ii. Arm B: Docetaxel 75mg/m2 IV q3w

Open 11/13/25

Mesothelioma (C45)

- 1.

Lung, Small Cell (C34)

- 1. Extensive Stage, Small Cell Lung Ca
 - a. **BMS CA245-0001 A Randomized, Double-Blind, Multicenter Phase 3 Trial**
 - i. Arm A: BMS-986489 420mg/nivolumab 360 mg IV+carboplatin+etoposide
 - ii. Arm B: atezolizumab 1200mg IV+carboplatin+etoposide
- 2. Previously Treated

Open 1/24/25

Breast, Adjuvant (C50)

- 1. NeoAdjuvant, Stage I, II, III
- 2. Adjuvant
 - a. **GS-US-595-6184 (ASCENT-05) A Randomized, Open-label, Phase 3 Study of Adjuvant Sacituzumab Govite**
 - i. Arm A: Sacituzumab Govitecan 10mg/kg IV Days 1 and 8 + Pembrolizumab 200mg IV Day 1 Q3W X 8 cycles
 - ii. Arm B: Treatment of Physician's Choice: Pembrolizumab 200mg IV Day 1 Q3 weeks X 8 cycles OR Pembrolizumab 200 mg IV Day 1 and Capecitabine 1000 mg/m2 PO BID D1-D14 Q3 Weeks X 8 cycles

Open 1/31/23

Breast, Her-2 Positive (C50)

- 1. NeoAdjuvant HER-2 Positive
- 2. Adjuvant, HER-2 Positive
- 3. Metastatic, HER-2 Positive, 1st Line
- 4. Metastatic, Previously Treated, Her-2 Positive

Breast, Advanced (C50)

- 1. Stage IV, Metastatic, 1st Line
 - a. **RGN WO45654 (INAVO123) A PHASE III, MULTICENTER, RANDOMIZED, DOUBLE-BLIND, PLACEBO-CONTROLLED**
 - i. Arm A: Inavolisib 9mg tablet PO QD days 1-28 of 28 day cycle + + letrozole 2.5mg PO QD + Investigator's choice - palbociclib 125mg PO QD D1-21 of 28 or ribociclib 600mg D1-21 of 28d cycle
 - ii. Arm B: Placebo tablet PO QD days 1-28 of 28 day cycle + + letrozole 2.5mg PO QD + Investigator's choice - palbociclib 125mg PO QD D1-21 of 28 or ribociclib 600mg D1-21 of 28d cycle
- 2. Stage IV, Metastatic, ER -
- 3. Stage IV, Metastatic, Previously Treated
- 4. Stage IV, Metastatic, Triple Negative Breast Cancer

Open 5/23/25

Breast, ER Positive (C50)

- 1. Adjuvant
- 2. Metastatic, First Line
 - a. **AZ D8532C00008 (SERAF-A-1) Phase IIb study to evaluate Camizestrant plus Ribociclib in ER-positi**
 - i. Camizestrant 75 mg PO QD for 28 days + Ribociclib 600 mg PO QD for 21 days of a 28 day cycle
- 3. Metastatic, Prior Hormonal Therapy

Pending 6/24/26

Melanoma (C43)

- 1. Adjuvant
- 2. Locally advanced or Metastatic
- 3. Previously Treated

Ovarian (C56)

- 1. Previously Treated

Bladder (C67)

- 1.
- 2. Urothelial

Renal (C64)

- 1. Adjuvant, Phase III
- 2. First Line Treatment, Advanced or Metastatic
- 3. Advanced, Metastatic
 - a. **MK 6482-034 A Phase 3, Randomized, Double-blind, Study of Belzutifan + Zanzalintin**
 - i. Arm 1: Belzutifan 120 mg qd+ Zanzalintinib 60 mg qd
 - ii. Arm 2: Belzutifan 120 mg qd+Placebo
- 4. Previously Treated

Pending 5/29/26
Accrual = 3

Prostate (C61)

- 1. Hormone Refractory, 1st Line
- 2. 1st line Metastatic
- 3. Metastatic, 2nd Line Prostate
- 4. Hormone Refractory; Bone Predominant 3rd Line or Greater

Sarcoma (C46-C49)

- 1. Sarcoma, Previously Treated

Adrenal (C74)

- 1. Previously Treated

Lymphoma (C82-C88)

- 1. Indolent, previously treated
- 2. Follicular, First Line
- 3. Indolent, Relapsed
- 4. Relapsed or Refractory; DLBCL, FL
- 5. Large Cell Lymphoma, Relapsed/Refractory
- 6. Relapsed and Refractory Follicular Lymphoma
 - a. **BMS CA073-1003 A Phase 3, Multicenter, Randomized, Open Label Study to Compare the Ef**
 - i. Arm A: R-Golca 0.4mg D1-14/48 days+Rituximab x5cycles followed by golcadomide monotherapy for up to 12 cycles of toral therapy
 - ii. Arm B: R-Len 20mg, Days 1-21, cycles 1-12,rituximab 375mg/m2, Days 1,8,15,and 22 in cycle 1; Day 1 in cycles 2-5
 - iii. Arm C: Inv. Choice for 6 cycles: R-CHOP or Rituximab-Bendamustine
- 7. Relapsed and Refractory Follicular or Marginal Zone Lymphoma
- 8. Diffuse Large Cell, First Line
- 9. Mantle Cell Lymphoma

Open 8/13/25

Accrual Goal = 3

CNS (C71)

- 1. Glioblastoma Multiforme

Leukemia (C91-C92)

- 1. CLL, First Line
- 2. CLL, Relapsed or Refractory
- 3. CML

Myeloma (C90)

- 1. 1st Line
- 2. Relapsed / Refractory 1-3 Lines
- 3. Relapsed / Refractory > or equal to 2 Lines

MDS (D46)

- 1. 1st Line
- 2. Observational Registry

Unknown Primary (199)

- 1. 1st Line

Other

- 1. Tissue Studies

- 2. Observational Studies

Multiple Sites

- 1. Phase I
- 2. EMD MS100070-0176 (EMR100070-001 Rollover)
- 3. Novartis Signature Series
- 4. Solid Tumors, Refractory
- 5. Solid Tumors, Advanced or Metastatic

Supportive Care

- 1. Bone Mets
- 2. Anemia
- 3. Thrombocytopenia

this list was last updated on: 7/23/26